



William J. Mesibov MD  
Linda H. Jacobs MD

Stuart J. Altman MD  
Marcia Rubinos MD

Fellows American Academy of Pediatrics  
575 Underhill Boulevard Syosset, New York 11791 516.921.2122 [www.kidfixer.com](http://www.kidfixer.com)

## Tics (Twitches)

Definition. Tics are:

-> Rapid, repeated muscle twitches (also called habit spasms), such as eye blinking, facial grimacing, forehead wrinkling, head jerking, or shoulder shrugging.

-> Usually last only 1 second.

-> Increase with stress.

-> Decrease with relaxation and disappear during sleep.

-> Occur in 20% of children and are three times more frequent in boys than in girls

-> Occur most often in children 5 to 10 years old.

### Causes

Most motor tics are due to inherited biochemical differences, not emotional problems. Flurries of tics, however, may indicate that your child is under pressure. All tics are involuntary, not deliberate. Children who have tics are usually normal, bright, and sensitive. Tics can be worsened by critical parents who nag or pressure a child.

### Expected Course

If tics are ignored, they usually disappear in 2 months to 1 year. If extra effort is made to help your child relax, they may improve more quickly. Even if the tics are not ignored and a child continues to feel stress or pressure, the tics usually improve or clear spontaneously during adolescence.

### How to Help Your Child with Tics

-> Help your child to relax in general. Make sure your child has free time and fun time every day. If your child is overscheduled with activities, try to lighten the commitments. If your child is unduly self-critical, praise him more and remind him to be a good friend to himself.

-> Identify and remove specific environmental stresses. Whenever your child has a flurry of tics, write in a diary the date, time, and preceding event. From this diary, you may be able to identify when your child feels pressure (Note: Your child should not know that you are keeping this

diary). In general, criticize your child less about grades, music lessons, sports, keeping his room clean, table manners, and so forth. Avoid stimulant medications (such as decongestants), which can lower the threshold for tics.

-> Ignore tics when they occur. When your child is having tics, don't call his attention to them. Reminders imply that they are bothering you. If your child becomes worried about the tics, then every time they occur, the child will react with tension rather than acceptance. The tension in turn will trigger more tics. Don't allow siblings to tease your child about the tics. Be sure that relatives, friends, and teachers also ignore the tics. When tics occur, people should focus on reducing any pressure they may be causing your child.

-> Don't talk about tics when they are not occurring. Stop all family conversation about tics. The less said about them, the less your child will be apprehensive of them. If your child brings up the subject, say something reassuring, such as, "Eventually your muscles will learn to relax and the tics will go away."

-> Avoid any punishment for tics. Some parents have the mistaken idea that tics are a bad habit that can be broken. This idea is absolutely false. If a child is made to practice "controlling tics" in front of a mirror, he will realize only that he cannot control them and they will become worse. Any facial exercises or massage should be discontinued because it only draws undue attention to the problem.

### Call Our Office During Regular Hours If:

- »The tics interfere with friendships or studies at school.
- »The tics involve sounds or words.
- »The tics become much more frequent.
- »The tics have lasted for more than a year.
- »You have other questions or concerns.



Scan this with your smart phone scanning app and you'll see the Kidfixer website.